

Topological Speed Limits to Network Synchronization

Marc Timme, Fred Wolf, and Theo Geisel

Max-Planck-Institut für Strömungsforschung, 37073 Göttingen, Germany

We study collective synchronization of pulse-coupled oscillators interacting on asymmetric random networks. We demonstrate that random matrix theory can be used to accurately predict the speed of synchronization in such networks in dependence on the dynamical and network parameters. Furthermore, we show that the speed of synchronization is limited by the network connectivity and stays finite, even if the coupling strength becomes infinite. In addition, our results indicate that synchrony is robust under structural perturbations of the network dynamics.

PACS numbers: 05.45.-a, 89.75.-k, 89.20.-a, 87.10.+e

Complex networks have attracted an enormous research interest in the recent past [1]. Most studies have focussed on the structures of a variety of systems such as the world wide web, email networks, genetic networks, and biological neural networks [1, 2]. An equally important task is to understand the *collective dynamics* on such networks. In particular, the question arises: how is the dynamics on a complex network influenced by its structure [2]?

Synchronization appears to be one of the simplest kinds of collective dynamics among coupled dynamical systems [3, 4]. It was found to be ubiquitous in artificial as well as natural networks as different as Josephson junction arrays [5] and biological neural networks [6]. To understand the dynamics of such networks, theoretical studies have emphasized systems consisting of simple units such as phase- and pulse-coupled limit-cycle oscillators [7, 8]. Yet, although most real-world networks often possess a complex connectivity structure, most studies of synchronization of coupled oscillators are either restricted to networks of globally coupled units and simple regular networks, or work in some mean field limit [7, 8]. Although exact results on synchronization in networks with a general structure have been obtained recently [9, 10, 11], it is still not well understood how the structure of a complex network affects dynamical features of synchronization.

In this Letter, we study the collective synchronization of pulse-coupled oscillators interacting on asymmetric random networks. We find that the speed of synchronization is restricted by the network connectivity and stays finite, even if the coupling strength becomes infinite. No such speed limit exists in large networks of globally coupled units. More generally, we show that the theory of random matrices can be used to successfully predict the speed of synchronization as a function of dynamical and network parameters. In addition, our results indicate that synchrony occurs robustly, i.e. persists under structural perturbation of the network dynamics.

We consider asymmetric random networks of N oscillators which interact by sending and receiving pulses [12]. The sets $\text{Pre}(i)$ of presynaptic oscillators that send pulses

to oscillator i specify the structure of such a network. For each oscillator i , the $k_i := |\text{Pre}(i)|$ presynaptic oscillators are drawn from the uniform distribution among all other oscillators $\{1, \dots, N\} \setminus \{i\}$.

A phase-like variable $\phi_i(t)$ specifies the state of each oscillator i at time t . In the absence of interactions, the dynamics of unit i is given by $d\phi_i/dt = 1$. When oscillator i reaches a threshold, $\phi_i(t) = 1$, its phase is reset to zero, $\phi_i(t^+) = 0$, and the oscillator is said to 'fire'. A pulse is sent to all postsynaptic oscillators $j \in \text{Post}(i)$ which receive this signal after a delay time τ . The incoming signal induces a phase jump $\phi_j((t+\tau)^+) := U^{-1}(U(\phi_j(t+\tau)) + \varepsilon_{ji})$ which depends on the instantaneous phase $\phi_j(t+\tau)$ of the postsynaptic oscillator and the coupling strength ε_{ji} which we take to be inhibitory (phase-retarding), $\varepsilon_{ji} \leq 0$. The phase dependence is determined by a twice continuously differentiable 'potential' function $U(\phi)$ that is assumed to be strictly increasing, $U'(\phi) > 0$, concave (down), $U''(\phi) < 0$, and normalized such that $U(0) = 0$, $U(1) = 1$ (cf. [12]). We focus on the specific form $U(\phi) = U_{\text{IF}}(\phi) = I(1 - e^{-T_{\text{IF}}\phi})$ that represents the integrate-and-fire oscillator defined by the differential equation $\dot{V} = I - V$ (and a threshold at $V = 1$). Here $I > 1$ is an external input and $T_{\text{IF}} = \log(I/(I-1))$ the intrinsic period of an oscillator. Other forms of $U(\phi)$ give qualitatively similar results. In such a network the synchronous state, $\phi_i(t) = \phi_0(t)$ for all i , exists if the coupling strengths are normalized such that $\sum_{j \in \text{Pre}(i)} \varepsilon_{ij} = \varepsilon$. Its period is given by $T = \tau + 1 - U^{-1}(U(\tau) + \varepsilon)$.

In numerical simulations of the network dynamics, we find that the synchronous state is always stable, independent of the parameters (cf. [10]). A sufficiently small perturbation $\delta(0) \equiv \delta = (\delta_1, \dots, \delta_N)^T$ of the phases, defined by $\phi_i(0) = \phi_0(0) + \delta_i$ asymptotically decays exponentially with time. Thus, denoting $\delta'(t) := \delta(t) - \lim_{s \rightarrow \infty} \delta(s)$, the distance $\Delta(n) := \max_i |\delta'_i(nT)| / \max_i |\delta'_i(0)|$ from the synchronous state behaves as

$$\Delta(n) \sim \exp(-n/\tau_{\text{syn}}) \quad (1)$$

defining a synchronization time τ_{syn} in units of the collective period T . The speed of synchronization τ_{syn}^{-1} strongly

depends on the parameters. For instance, as might be expected, synchronization is faster for stronger coupling. Surprisingly, however, we find that synchronization cannot be faster than an upper bound even if the coupling strength becomes arbitrarily large (cf. Fig. 1).




FIG. 1: Asymptotic synchronization time in a random network ($N = 1024$, $k_i \equiv k = 32$, $I = 1.1$, $\tau = 0.05$, $\varepsilon_{ij} = \varepsilon/k$ for $j \in \text{Post}(i)$). The inset shows the distance Δ of a perturbation δ from the synchronous state versus the number of periods n ($\varepsilon = -0.4$). Its slope yields the synchronization time τ_{syn} shown in the main panel as a function of coupling strength $|\varepsilon|$. Simulation data ($\circ$), theoretical prediction (—) derived in this Letter, its infinite coupling strength asymptote (---).

To understand how the speed of synchronization depends on the dynamical and network parameters, we analyze the linear stability of the synchronous state. Following [10] we obtain a nonlinear stroboscopic map $\delta(T) = F(\delta)$ for the perturbations, the linearization of which reads

$$\delta(T) \doteq A\delta \quad (2)$$

where the elements of the stability matrix A are given by $A_{ij} = p_{i,n} - p_{i,n-1}$ if $j = j_n \in \text{Pre}(i)$, $A_{ii} = p_{i,0}$, and $A_{ij} = 0$ otherwise. Here j_n identifies the n^{th} pulse received during this cycle by oscillator i and $p_{i,n} := U'(U^{-1}(U(\tau) + \sum_{m=1}^n \varepsilon_{ij_m})) / U'(U^{-1}(U(\tau) + \varepsilon))$ for $n \in \{0, 1, \dots, k_i\}$. For $U(\phi) = U_{\text{IF}}(\phi)$ and coupling strengths $\varepsilon_{ij} = \varepsilon/k_i$ for $j \in \text{Pre}(i)$ the matrix elements reduce to [13]

$$A_{ij} = \begin{cases} \frac{1-A_0}{k_i} & \text{if } j \in \text{Pre}(i) \\ A_0 & \text{if } j = i \\ 0 & \text{if } j \notin \text{Pre}(i) \cup \{i\} \end{cases} \quad (3)$$

where

$$A_0 = \frac{Ie^{-\tau T_{\text{IF}}}}{Ie^{-\tau T_{\text{IF}}} - \varepsilon} > 0. \quad (4)$$

Obviously, A constitutes a row-stochastic matrix, i.e. $\sum_j A_{ij} = 1$ for all i . Thus A has one trivial eigenvalue $\lambda_1 = 1$ associated with the eigenvector $\mathbf{v}_1 = (1, 1, \dots, 1)^T$

representing a uniform phase-shift and thus reflecting time-translation invariance. Furthermore, the Gershgorin theorem [15] implies that all eigenvalues are located inside a disk of radius $r_G = 1 - A_0$ centered at A_0 , such that in particular $|\lambda_i| \leq 1$ and the synchronous state is (at least neutrally) stable. For simplicity, we consider networks of homogenous random connectivity, $k_i = k$ for all i , in the following.

We numerically determined the eigenvalues of different stability matrices A for various network sizes $N \in \{2^6, \dots, 2^{14}\}$, in-degrees $k \in \{2, \dots, 2^8\}$, and dynamical parameters ε , τ , and I such that $0 < A_0 < 1$. In general, we find that for sufficiently large k and N the non-trivial eigenvalues resemble a disk in the complex plane that is centered at about A_0 and has a radius r that is smaller than the upper bound given by the Gershgorin theorem, $r < 1 - A_0$. Examples are shown in Fig. 2.




Figure 2: Distribution of eigenvalues λ_i of two stability matrices A in the complex plane ($I = 1.1$, $\varepsilon = -0.2$, $\tau = 0.05 \Rightarrow A_0 \approx 0.83$; $k = 8$) for networks of (a) $N = 32$, (b) $N = 512$ oscillators. For large networks, the non-trivial eigenvalues seem to be distributed uniformly on a disk in the complex plane. The prediction from random matrix theory (Eq. (10)) is indicated by a circle. The arc through the trivial eigenvalue $\lambda_1 = 1$ is a sector of the unit circle.

This eigenvalue distribution is reminiscent of the “circle law” of random matrix theory [16]: Gaussian asymmetric random matrices, having a distribution of matrix elements

$$p_{\text{Gauss}}(J_{ij}) = N^{\frac{1}{2}} (2\pi r^2)^{-\frac{1}{2}} \exp\left(-\frac{N J_{ij}^2}{2r^2}\right) \quad (5)$$

with independent J_{ij} and J_{ji} , also exhibit eigenvalue distributions

$$\rho_{\text{Gauss}}^a(\lambda) = \begin{cases} (\pi r^2)^{-1} & \text{if } |\lambda| \leq r \\ 0 & \text{otherwise} \end{cases} \quad (6)$$

for $N \rightarrow \infty$ that are uniform in a disk in the complex plane [16]. The radius r of the disk is given by

$$r = N^{\frac{1}{2}} \sigma \quad (7)$$

where $\sigma^2 = \langle J_{ij}^2 \rangle$ is the variance of the matrix elements.

Interestingly, we find that the radii of the eigenvalue distributions of the above stability matrices (3) well

agree with the radii obtained from Eq. (7) if $\langle J_{ij}^2 \rangle$ is replaced by the variance of the elements of the stability matrices shifted such that they also exhibit a zero average eigenvalue. To directly compare the eigenvalues of the stability matrices, which have average eigenvalue $[\lambda_i] := \frac{1}{N} \sum_{i=1}^N \lambda_i = A_0$ to those of the Gaussian ensemble, we transform $A'_{ij} = A_{ij} - \delta_{ij} A_0$ shifting the average eigenvalues to $[\lambda'_i] = 0$. Here δ_{ij} denotes the Kronecker delta, $\delta_{ij} = 1$ if $i = j$ and $\delta_{ij} = 0$ if $i \neq j$. For the variance of A' we obtain

$$\sigma_{A'}^2 = [A_{ij}'^2] - [A'_{ij}]^2 \quad (8)$$

$$= \frac{1}{N} \left(\sum_{\substack{j=1 \\ j \neq i}}^N A_{ij}'^2 - \frac{(1 - A_0)^2}{N} \right). \quad (9)$$

For identical non-zero coupling strengths, the off-diagonal sum is exactly equal to $\sum_{j=1, j \neq i}^N A_{ij}'^2 = (1 - A_0)^2/k$ such that, using (7), we obtain the random matrix theory prediction

$$r_{\text{RMT}} = N^{\frac{1}{2}} \sigma_{A'} = (1 - A_0) \left(\frac{1}{k} - \frac{1}{N} \right)^{\frac{1}{2}}. \quad (10)$$

for the radius r of the disk of eigenvalues of the stability matrices A [17].

We verified this scaling law (10) for various dynamical parameters A_0 (determined by different I , ε , and τ), network sizes N , and in-degrees k and found excellent agreement with numerically determined eigenvalue distributions, see, e.g., Fig. 2. To quantify the accuracy of the prediction (10) we numerically estimated the radius of the distribution of the non-trivial eigenvalues of A for various N , k as well as A_0 . Results from two different estimators are shown in Fig. 3. The real part estimator $r_{\text{Re}} := \frac{1}{2} (\max_{i \neq 1} \text{Re}(\lambda_i) - \min_{i \neq 1} \text{Re}(\lambda_i))$ estimates the radius from the maximum spread of eigenvalues parallel to the real axis. Typically, r_{Re} should give an estimate that is slightly inaccurate because it is based on two eigenvalues only. This is circumvented by the average estimator $r_{\text{av}} := \frac{3}{2} \frac{1}{N-1} \sum_{i=2}^N |\lambda_i - (A_0 - (1 - A_0)N^{-1})|$ that estimates the radius r of a circle from the average distance $\langle d \rangle$ of eigenvalues from its center, because $\langle d \rangle = \int_0^{2\pi} \int_0^r r'^2 \rho(r') dr' d\varphi = \frac{2}{3}r$ if we assume a uniform density $\rho(r')$ according to (6). Here we take the center of the disk to be the average $\langle \lambda_i \rangle_{i \geq 2} = A_0 - (1 - A_0)N^{-1} + \mathcal{O}(N^{-2})$ of the non-trivial eigenvalues. Varying k at fixed N as well as N at fixed k yields excellent agreement between the numerical data and the theoretical predictions for sufficiently large N and k (Fig. 3). Varying the coupling strength $|\varepsilon|$ and thus A_0 yields equally good agreement (cf. Fig. 1).

The radius (10) implies a prediction for the synchronization time (see (1))

$$\tau_{\text{syn}} = -1/\ln(A_0 + r_{\text{RMT}}), \quad (11)$$




Figure 3: Scaling of the radius r of the disk of non-trivial eigenvalues. Main panel displays the radius r as a function of network size N for fixed $k = 32$. Symbols display r_{Re} ($\times$) and r_{av} ($\circ$). Inset displays r as a function of k for fixed $N = 1024$. Dots display numerical data of r_{av} . In the main panel and the inset, lines are the theoretical estimate r_{RMT} (Eq. (10)) Other parameters as in Fig. 2.

in terms of the (in modulus) largest non-trivial eigenvalue λ_m . With increasing coupling strength $|\varepsilon|$, the synchronization time decreases. However, the speed of synchronization τ_{syn}^{-1} is bounded by a finite speed for arbitrary large $|\varepsilon|$: Even if $|\varepsilon| \gg 1$ and thus $A_0 \ll 1$, the largest non-trivial eigenvalue asymptotically becomes $\lambda_m \approx k^{-1/2}$ for sufficiently large N . Thus the shortest synchronization time

$$\tau_{\text{syn}}^{|\varepsilon| \rightarrow \infty} = \frac{2}{\ln k} \quad (12)$$

is limited by the network connectivity (cf. the asymptote in Fig. 1). This means that even for arbitrary strong interactions, the speed of synchronization stays finite. Furthermore, at fixed k , the synchronization time also cannot exceed a certain maximum, even if the network size N becomes extremely large (cf. Fig. 3). This bound $\tau_{\text{syn}}^{N \rightarrow \infty}$ is determined by the asymptotic radius $r_\infty := \lim_{N \rightarrow \infty} r_{\text{RMT}} = (1 - A_0)k^{-1/2}$. Moreover, because eigenvalues change continuously with a structural perturbation to the system's dynamics, the existence of a gap $g := 1 - (A_0 + r_\infty) > 0$ indicates that no eigenvalue crosses the unit circle for sufficiently small structural perturbations. Thus stable synchrony is not restricted to the specific model considered here, but persists in systems obtained by structural perturbations of the dynamics.

The above results show, that the distribution of eigenvalues of a *sparse* stability matrix with deterministic non-zero entries at certain random positions is well described by the eigenvalue distribution of the *Gaussian* ensemble, which consists of fully occupied matrices with purely random entries. This sparse-Gaussian coincidence for asymmetric matrices is similar to that of symmetric random matrices for large k : Gaussian symmetric matrices exhibit an eigenvalue distribution $\rho_{\text{Gauss}}^s(\lambda)$, the

Wigner semicircle law [20]. Sparse symmetric matrices [18] exhibit an eigenvalue distribution $\rho_{\text{sparse}}^s(\lambda)$ that is different from the semicircle law but approaches it in the limit $k \rightarrow \infty$. For $k \gg 1$ the distribution of eigenvalues of sparse asymmetric random matrices ρ_{sparse}^a appear to be well approximated by the eigenvalue distribution of Gaussian asymmetric matrices, $\rho_{\text{sparse}}^a(\lambda) \approx \rho_{\text{Gauss}}^a(\lambda)$. Our results indicate, that this is true even for moderate $k \approx 10$.

Further investigations of eigenvalue distributions for small-world networks show that with decreasing randomness the speed of synchronization decreases (the second largest non-trivial eigenvalue increases) such that completely random networks synchronize faster than small-world networks, at least asymptotically.

In conclusion, we have derived accurate analytical predictions for the (asymptotic) speed of synchronization in asymmetric random networks of oscillators in dependence of the dynamical parameters ε , τ , I , as well as the network parameters N and k . Even the scaling with network size N , artificially introduced via the variance (9) of finite matrices, is also accurately reproduced (see e.g. Fig. 3). As a particular application, we explained the intriguing phenomenon, that the speed of synchronization stays finite, even if the coupling strengths become infinite. It turned out that the speed is restricted by the network connectivity. In addition, at fixed parameters the synchronization time does not increase above a finite threshold if the network becomes very large. Furthermore, the existence of a gap between the non-trivial eigenvalues and the unit circle indicates that stable synchrony is a robust form of collective dynamics in a large class of systems.

Random matrix theory has previously been applied to various physical systems that exhibit certain symmetries – such as time-reversal symmetry – but an otherwise unknown structure. For instance, correlations of energy levels in nuclear physics and quantum mechanical properties of classically chaotic systems have been successfully predicted (see Ref. [19] for a recent review). Our results demonstrate that random matrix theory also is an appropriate tool for analyzing synchronization in random networks of dynamical units. Possible lines for future applications may include synchronization phenomena of pulse- and phase-coupled units as well as of chaotic dynamical systems (cf. also [21]). The limits of synchronization speed predicted in this Letter, are expected to occur in those systems, too. More generally, other equilibration processes and the dynamics in more structured topologies such as small-world networks may be analytically investigated using statistical spectral properties of the respective operators, too.

We thank T. Kottos, P. Müller, H. Sompolinsky, M. Weigt, and A. Zippelius for useful discussions.

-
- [1] S. Bornholdt and H. G. Schuster (eds.), *Handbook of Graphs and Networks*, (Wiley-VCH, Weinheim, 2002); R. Albert and A.-L. Barabási, *Rev. Mod. Phys.* **74**, 47 (2002); S. N. Dorogovtsev and J. F. F. Mendes, *Adv. Phys.* **51**, 1079 (2002); M. E. J. Newman, *SIAM Review* **45**, 167 (2003).
 - [2] S. H. Strogatz, *Nature* **410**, 268 (2001).
 - [3] S. Nadis, *Nature* **421**, 780 (2003).
 - [4] A. Pikovsky, M. Rosenblum, and J. Kurths, *Synchronization: A Universal Concept in Nonlinear Science* (Cambridge University Press, Cambridge, England, 2003).
 - [5] K. Wiesenfeld, P. Colet, and S. H. Strogatz, *Phys. Rev. Lett.* **76**, 404 (1996).
 - [6] C. M. Gray, P. König, A. K. Engel, and W. Singer, *Nature* **338**, 334 (1989); R. Eckhorn *et al.* *Biol. Cybern.* **60** 121 (1988); W. Singer, *Neuron* **24**, 49 and 111 (1999).
 - [7] A. T. Winfree, *The Geometry of Biological Time* (Springer, New York, 1980); Y. Kuramoto, *Chemical Oscillations, Waves, and Turbulence* (Springer, Berlin, 1984); for a recent review on the Kuramoto model see S. H. Strogatz, *Physica D* **143**, 1 (2000).
 - [8] A. V. M. Herz and J. J. Hopfield, *Phys. Rev. Lett.* **75**, 1222 (1995); Á. Corral, C. J. Pérez, A. Díaz-Guilera, and A. Arenas, *Phys. Rev. Lett.* **75**, 3697 (1995); W. Gerstner, J. D. Cowan, and J. L. van Hemmen, *Neural Comp.* **8**, 1653 (1996); P. C. Bressloff, S. Coombes, and B. de Souza, *Phys. Rev. Lett.* **79**, 2791 (1997); C. van Vreeswijk, *Phys. Rev. Lett.* **84**, 5110 (2000); D. Hansel and G. Mato, *Phys. Rev. Lett.* **86**, 4175 (2001).
 - [9] M. Barahona and L. M. Pecora, *Phys. Rev. Lett.* **89**, 054101 (2002).
 - [10] M. Timme, F. Wolf, and T. Geisel, *Phys. Rev. Lett.* **89**, 258701 (2002).
 - [11] M. G. Earl and S. H. Strogatz, *Phys. Rev. E* **67**, 036204 (2003).
 - [12] R. E. Mirollo and S. H. Strogatz, *SIAM J. Appl. Math.* **50**, 1645 (1990); U. Ernst, K. Pawelzik, and T. Geisel, *Phys. Rev. Lett.* **74**, 1570 (1995); M. Timme, F. Wolf, and T. Geisel, *Chaos* **13**, 377 (2003).
 - [13] In general, the stability matrix depends on the order of incoming signals. In this Letter, however, we focus on a subclass of models, for which this complication does not arise because the stability matrix becomes unique [14].
 - [14] M. Timme, *Collective Dynamics in Networks of Pulse-Coupled Oscillators*, Doctoral Thesis, University of Göttingen, 2002.
 - [15] J. Stoer and R. Burlisch, *Introduction to Numerical Analysis* (Springer, Berlin, 1992).
 - [16] V. L. Girko, *Theory Probab. Appl.* **29**, 694 (1985). H. J. Sommers, A. Crisanti, H. Sompolinsky, and Y. Stein, *Phys. Rev. Lett.* **60**, 1895 (1988).
 - [17] Here $k = k_i$ for all i such that the estimate for the radius (10) is exact. If the random network is constructed by choosing every connection independently with probability p , it becomes an approximation if k^{-1} is replaced by $[k_i^{-1}]_i$.
 - [18] G. J. Rodgers and A. J. Bray, *Phys. Rev. B* **37** 3557 (1988).
 - [19] P. J. Forrester, N. C. Snaith, and J. J. M. Verbaarschot, *J. Phys. A* **36**, R1 (2003).
 - [20] M. L. Mehta, *Random Matrices* (Academic Press, New

York, 1991).

[21] P. M. Gade, Phys. Rev. E **54**, 64 (1996).